



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification⁶ : A61K 38/04, 38/05, 31/12	A1	(11) International Publication Number: WO 98/13061 (43) International Publication Date: 2 April 1998 (02.04.98)
(21) International Application Number: PCT/US97/17013 (22) International Filing Date: 23 September 1997 (23.09.97) (30) Priority Data: 08/719,042 24 September 1996 (24.09.96) US (71) Applicant (for all designated States except US): CV THERAPEUTICS, INC. [US/US]; 3172 Porter Drive, Palo Alto, CA 94304 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): LUM, Robert, T. [US/US]; 781 Barron Avenue, Palo Alto, CA 94306 (US). SCHOW, Steven, R. [US/US]; 204 Mendocino Way, Redwood City, CA 94065 (US). JOLY, Alison [US/US]; 3205 Monterey Street, San Mateo, CA 94403 (US). KERWAR, Suresh [US/US]; 90 Edgewood Road, Westchester, NY 10562 (US). NELSON, Marek, G. [US/US]; 2690 Park Way, Sunol, CA 94586 (US). WICK, Michael, M. [US/US]; 16 Beresford Road, Chestnut Hill, MA 02146 (US). (74) Agent: HUGHES, A., Blair, McDonnell Boehnen Hulbert & Berghoff, 300 South Wacker Drive, Chicago, IL 60606 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TS, TT, UA, UG, US, UZ, VN, ARIPO patent (GH, GL, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: INHIBITION OF 26S AND 20S PROTEASOME BY INDANONES (57) Abstract This invention is a method for inhibiting cell proliferation using indanones.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

**TITLE: INHIBITION OF 26S AND 20S
PROTEASOME BY INDANONES**

BACKGROUND OF THE INVENTION

1. Field of Invention

10 This invention is a method for inhibiting cell proliferation using a class of indanone compositions never before considered for that purpose. As inhibitors of cell proliferation, the compositions are useful in the treatment of cancer, cardiovascular disease, e.g. restenosis, host graft rejection, gout, and other proliferative disorders as well as being potential therapeutics for autoimmune diseases, such as rheumatoid arthritis, lupus, type I diabetes,
15 multiple sclerosis and similar disorders and diseases.

2. Description of the Art

The multicatalytic proteinase or proteasome is a highly conserved cellular structure that is responsible for the ATP-dependent proteolysis of most cellular proteins. The 20S (700-kDa) proteasome contains at least five distinct proteolytic activities that have a new type
20 of mechanism involving a threonine residue at the active site (Coux, O., Tanaka, K. and Goldberg, A. 1996 Ann. Rev. Biochem. 65:801-47).

The 20S proteasome has been crystallized from the archaebacteria *Thermoplasma acidophilum* (Lowe, J., Stock, D., Jap, B., Zwickl, P., Bauminster, W., and Huber, R. 1995 Science 268:533-539). The archaebacterial 20S proteasome contains fourteen copies of two
25 distinct types of subunits, α and β , which form a cylindrical structure consisting of four stacked rings. The top and bottom rings contain seven α subunits each whilst the inner rings

contain seven β subunits. A pore extends through the middle of the structure that contains the proteolytic active sites and proteins destined for degradation pass through this channel. The eukaryotic 20S proteasome is more complex than that of the archaebacteria because the number of distinct subunits has increased during evolution, however, the subunits can still be

5 classed according to the α and β nomenclature of archaebacteria according to their homology. Thus the quaternary structure of the eukaryotic complex is similar to that of the archaebacteria being comprised of two α and two β rings. However, unlike the archaebacterial proteasome that primarily exhibits chymotrypsin-like proteolytic activity (Dahlmann,B., Kopp,F., Kuehn,L., Nidel,B., Pfeifer,G. 1989 FEBS Lett. 251:125-131,

10 Seemuller, E., Lupas,A., Zuhl, F., Zwickl, P and Baumeister, W. 1995 FEBS Lett. 359:173; and Lowe,J., Stock,D., Jap,B., Zwickl,P., Bauminster, W. and Huber,R. 1995 Science 268:533-539). The eukaryotic proteasome contains at least five identifiable protease activities. These are termed chymotrypsin-like, trypsin-like and peptidylglutamyl-peptide hydrolyzing. Two other activities have also been described, one exhibiting a preference for

15 cleavage of peptide bonds on the carboxyl side of branched chain amino acids and the other toward bonds between small neutral amino acids. (Orlowski, M. 1990 Biochemistry 29: 10289-10297).

Although the 20S proteasome contains the proteolytic core, it cannot degrade proteins *in vivo* unless it is complexed with a 19S cap at either end of its structure, which itself

20 contains multiple ATPase activities. This larger structure is known as the 26S proteasome and rapidly degrades proteins that have been targeted for degradation by the addition of multiple molecules of the 8.5-kDa polypeptide, ubiquitin.

The first step towards the ubiquitination of a protein proceeds by activation of a ubiquitin molecule at its carboxyl terminal glycine residue by addition of ATP that creates a high energy thioester intermediate. This step is catalyzed by the ubiquitin-activating enzyme, E1. Ubiquitin is then transferred to the active cysteine residue of a ubiquitin-conjugating enzyme, E2. E2 enzymes attach ubiquitin to the E-amino groups of lysine residues on the substrate protein that is destined to be degraded. This process, in some cases also requires a ubiquitin ligase, E3. Repeated conjugation of ubiquitin to lysine residues of formerly bound ubiquitin moieties leads to the formation of multi-ubiquitin chains and creates a scaffold of ubiquitin around the substrate protein. Multi-ubiquitinated substrate proteins are recognized by the 26S proteasome and are degraded and the multi-ubiquitin chains are released from the complex and ubiquitin is recycled.

What causes a protein to become ubiquitinated and thus degraded is still under investigation. Clearly this must be a highly regulated series of events since the critical timing of specific protein degradation is crucial for many cell cycle functions. Several signals have been proposed which largely center upon internal structural sequences within the substrate itself. One such proposal is the "N-end rule" in which the amino terminal residue of a protein determines its half life. Other proteins such as the cyclins contain a short sequence of highly conserved amino acids termed the "destruction box" that are apparently necessary for degradation. Furthermore "PEST" sequences, which consist of regions rich in proline, aspartate, glutamate, serine and threonine also seem to act as degradation signals. It is thought that such internal sequences act as recognition elements between the protein substrate and its specific ubiquitination machinery.

Two types of inhibitors that inhibit the proteolytic activity of the proteasome, have

been described. Certain peptide aldehydes have been reported to inhibit the chymotrypsin-like activity associated with the proteasome (Vinitsky,A., Michaud,C., Powers,J. and Orlowski,M. 1992 Biochemistry 31:9421-9428; Tsubuki,S., Hiroshi,K., Saito,Y., Miyashita,N., Inomata,M., and Kawashima,S., 1993 Biochem.Biophys.Res.Comm. 196:1195-1201; Rock,K.I., Gramm,C., Rothstein,L., Clark,K., Stein,R., Dick,L., Hwang,D., and Goldberg,A.L. 1994 Cell 78:761-771). These are N-acetyl-L-leuciny-L-leucinal-L-norleucinal (ALLN) and a closely related compound, N-acetyl-L-leuciny-L-leuciny-methional (LLM) with a K_i 's of 0.14 μ M. The most potent inhibitor of this type is a structurally related compound, N-carbobenzoxyl-L-leuciny-L-leuciny-L-norvalinal (MG 115) which exhibits a K_i of 0.021 μ M. Although these peptide aldehydes are most effective against the chymotrypsin-like proteolytic activity of the proteasomes, careful studies have shown that they are non-specific protease inhibitors. More recent reports have described series of potent dipeptide inhibitors that have IC_{50} values in the 10-100 nM range *in vitro* (Iqbal,M., Chatterjee,S., Kauer,J.C., Das,M., Messina,P., Freed,B., Biazzo,W., and Siman,R. 1995 J.Med.Chem. 38:2276-2277) and a series of similarly potent *in vitro* inhibitors from α -ketocarbonyl and boronic ester derived dipeptides (Iqbal, M., Chatterjee, S., Kauer, J. C., Mallamo, J. P., Messina,P.A., Reiboldt,A., and Siman,R. 1996 Bioorg. Med.Chem. Lett 6:287-290).

Another report describes a class of compounds that exhibit specificity in inhibiting proteasome activity (Fenteany,G., Standaert,R.F., Lane,W.S., Choi,S., Corey,E.J., and Schreiber,S.L. 1995 Science 268:726-731). Lactacystin is a *Streptomyces* metabolite that specifically inhibits the proteolytic activity of the proteasome complex. This molecule was originally discovered for its ability to induce neurite outgrowth in a neuroblastoma cell line

(Omura et al., 1991 J.Antibiot. 44:113) later it was shown to inhibit the proliferation of several cell types (Fenteany et al 1994 Proc.Nat'l. Acad.Sci. USA 91: 3358). By using radiolabelled lactacystin, binding studies by (Fenteany et al., 1995 Science 268:726-731) have identified the binding site and the mechanism of action. These studies have shown that
5 lactacystin binds irreversibly to a threonine residue located at the amino terminus of the β subunit of the proteasomes. A series of analogues based upon the structure of lactacystin were also investigated (Fenteany et al., 1995 Science 268:726-731). These studies indicated that the β -lactone structure was essential for its inhibitory activity.

It is now well established that the proteasome is a major extralysosomal proteolytic
10 system which is involved in the degradative pathways resulting in numerous and diverse cellular functions such as cell division, antigen processing and the degradation of short lived regulatory proteins such as transcription factors, oncogene products and cyclins (reviewed in Ciechanover,A. 1994 Cell 79:13-21). The primary function of the proteasome is to catalyze the proteolysis of proteins into small peptides. However, it has also been demonstrated that
15 the ubiquitin-proteasome pathway can catalyze the regulated proteolytic processing of a large inactive precursor into an active protein. The best documented case of this involves the activation of the transcription factor NF- κ B (Palombella, V.J., Rando, O.J., Goldberg,A.L., and Maniatis,T. 1994 Cell 78:773-785). The active form of NF- κ B is a heterodimer consisting of a p65 and a p50 subunit. The latter is present in the cytosol of the cell in an
20 inactive precursor form, namely p105, the 105-kDa polypeptide precursor of p50. The proteolytic processing of p105 to generate p50 occurs via the ubiquitin-proteasome pathway. Additionally, processed p50 and p65 is maintained in the cytosol as an inactive complex with the inhibitory protein I κ B. Inflammatory signals activate NF- κ B by initiating the signalling

pathway for the complete degradation of I κ B, and also stimulate the processing of p105 into p50. Thus two proteolytic events, both governed by the ubiquitin-proteasome pathway, are required for signal induced activation of NF- κ B. What causes the termination of proteolysis of p105 following generation of p50 is not known but it has been proposed that the
5 conformation of p50 may render it resistant to further processing and cause it to dissociate from the 26S complex.

The fact that the proteasome plays a critical event in the activation of NF- κ B could be exploited clinically by the use of inhibitors directed towards proteasome proteolysis. In certain diseases the normal function of active NF- κ B can be detrimental to human health as
10 observed in inflammatory responses following bacterial, fungal or viral infection. Thus inhibitors of NF- κ B activation, due to their ability to prevent secretion of cytokines, may have potential utility in the treatment of ARDS (acute respiratory distress syndrome) and AIDS. Since activation of NF- κ B is also essential for angiogenesis, proteasome inhibitors may have utility in the treatment of those diseases associated with abnormal
15 neovascularization.

p53 was first described as an oncoprotein but has since been shown to be involved in many cellular processes (reviewed by Ko, L.J. and Proves, C. 1996 *Genes Dev.* 10, 1054-1072). p53 has been shown to induce apoptosis in several haematopoietic cell lines (Oren, M., 1994 *Semin. Cancer Biol.* 5, 221-227) through the action of many different stimuli
20 including DNA damage, viral infection and the removal of growth factors. However, it is important to note that apoptosis can be induced in a p53-independent manner for example by the action of glucocorticoids. Induction of p53 leads to cell growth arrest in the G1 phase of the cell cycle as well as cell death by apoptosis. Both of these functions allow p53 to control

DNA damage thereby reducing the propagation of DNA mutations when cells divide. p53 arrests cells at G1 by inducing the cyclin-dependent kinase inhibitor, p21, which in turn causes an accumulation of the hypophosphorylated form of the retinoblastoma gene product. It is thought that p53 acts as a check point in the cell following DNA damage, it first causes an arrest in cell division and apoptosis. p53 degradation is known to be via the ubiquitin-proteasome pathway and disrupting p53 degradation is a possible mode of inducing apoptosis. Another potential utility of proteasome inhibitors may be in the treatment of diseases that result from abnormal cell proliferation.

It is well documented that the ubiquitin-proteasome pathway is critical for the regulated destruction of cyclins that govern the exit from mitosis and allow cells to progress into the next phase of the cell cycle. Thus inhibiting degradation of cyclins by using proteasome inhibitors causes growth arrest. Therefore another potential utility of proteasome inhibitors is their use in the treatment of diseases that result from an accelerated cell division. These include cancer, cardiovascular diseases such as myocarditis, restenosis following angioplasty, renal diseases such as lupus, polycystic kidney disease, fungal infections, dermatological diseases such as psoriasis, abnormal wound healing, keloids, immunological diseases such as autoimmunity, asthma, and allergy, acute and delayed hypersensitivity, graft versus host disease, transplant rejection and neuroimmunological diseases such as multiple sclerosis and acute disseminated encephalomyelitis.

SUMMARY OF THE INVENTION


It is an object of this invention to provide a method for inhibiting cell proliferation in mammals that uses a therapeutically effective amount of a composition heretofore unknown
5 for its cell proliferative inhibition properties.

It is an object of this invention to provide a method for the effective treatment of diseases that result from accelerated cell division.

It is another object of this invention to provide a method for the treatment of proliferative diseases that operates by inhibiting the degradation of proteasome inhibitors.

10 It is another object of this invention to use a therapeutically effective amount of the composition to inhibit cell proliferative disorders in humans.

In one embodiment, this invention is a method for inhibiting cell proliferation in mammals comprising administering to the mammal a therapeutically effective amount of a compound having the formula:




20

In the compound, R₁-R₄ are each individually selected from the group including hydrogen, halogen, hydroxyl, thiol, lower alkyl, substituted lower alkyl, alkenyl, alkynyl, alkylalkenyl, alkyl alkynyl, alkoxy, alkylthio, acyl, aryloxy, amino, amido, carboxyl,


aryl, substituted aryl, heterocycle, heteroaryl, substituted heterocycle, heteroalkyl, cycloalkyl, substituted cycloalkyl, alkylcycloalkyl, alkylcycloheteroalkyl, nitro, or cyano.

R_4, R_5 are each individually selected from the group of compounds including hydrogen, halogen, hydroxyl, thiol, oxo, lower alkyl, substituted lower alkyl, alkenyl, alkynyl, alkylalkenyl, alkyl alkynyl, alkoxy, alkylthio, acyl, aryloxy, amino, amido, carboxyl, aryl, substituted aryl, heterocycle, heteroaryl, substituted heterocycle, heteroalkyl, cycloalkyl, substituted cycloalkyl, alkylcycloalkyl, alkylcycloheteroalkyl, nitro, or cyano;

X is selected from the group of compounds including hydrogen, $-D_1$, $-D_2$, $-E$, $-D_1-E$, $-D_2-E$, $-D_1-D_2$ or a compound having the formula:



wherein D_1 and D_2 are each individually selected from the group of compounds including a compound having the formula:




20

or hydrogen, halogen, hydroxyl, thiol, lower alkyl, substituted lower alkyl, alkynyl, alkyl alkenyl, alkyl alkynyl, alkoxy, alkylthio, acyl, aryloxy, amino, amido, carboxyl, aryl, substituted aryl, heterocycle, heteroaryl, substituted heterocyclic, heteroalkyl,

cycloalkyl, substituted cycloalkyl, alkylcycloalkyl, or alkylcycloheteroalkyl;

wherein E is selected from the group of compounds including:



or hydrogen, halogen, hydroxyl, thiol, lower alkyl, substituted lower alkyl, alkynyl, alkylalkenyl, alkyl alkynyl, alkoxy, alkylthio, acyl, aryloxy, alkylthio, acyl, aryloxy, amino, amido, carboxyl, aryl, substituted aryl, heterocycle, heteroaryl, substituted
10 heterocycle, heteroalkyl, cycloalkyl, substituted cycloalkyl, alkylcycloalkyl, or alkylcycloheteroalkyl.

When D₁, D₂ and/or E are selected from compounds including substituents R₁₀-R₁₄, J₁ and J₂, then R₁₀-R₁₄ are each individually selected from the group of compounds including hydrogen, halogen, hydroxyl, thiol, oxo, lower alkyl, substituted lower alkyl, alkenyl, alkynyl, alkylalkenyl, alkyl alkynyl, alkoxy, alkylthio, acyl, aryloxy, amino, amido, carboxyl, aryl, substituted aryl, heterocycle, heteroaryl, substituted heterocycle, heteroalkyl, cycloalkyl, substituted cycloalkyl, alkylcycloalkyl, alkylcycloheteroalkyl, nitro, or cyano and J₁ and J₂ are each individually selected from the group of compounds including N-R₁₅, CR₁₆, R₁₇, O, S-(O)₀₋₂, P-(O)₀₋₃, wherein R₁₅-R₁₇ may each be individually selected from the group of compounds including hydrogen, halogen, hydroxyl, oxo, thiol, lower alkyl, substituted lower alkyl, alkynyl, alkylalkenyl, alkyl alkynyl, alkoxy, alkylthio, acyl, aryloxy, amino, amido, carboxyl, aryl, substituted aryl, heterocyclic, heteroaryl, substituted heterocycle, heteroalkyl, cycloalkyl, substituted cycloalkyl,

alkylcycloalkyl, alkylcycloheteroalkyl, or cyano.

The compositions are useful, when administered in therapeutic amounts, for treating mammals, and preferably for treating humans suffering from cell proliferative disorders, infectious diseases, and immunological diseases.

DESCRIPTION OF THE FIGURES


In the Figures, Figure 1 is a Western Blot immunoreactivity assay using an anti-I κ B α antibody of RAW cell extract which had been treated with compounds 173 and 187 which are described in Tables 1 and 2;

5 Figure 2 is a Western Blot immunoreactivity assay towards an anti-P50 antibody of RAW cell extracts which had been treated with compound 187 which is described in Tables 1 and 2 prior to exposure to its LPS; and

Figure 3 is a gel mobility shift assay using nuclear extract prepared from RAW cells that had been pretreated compound 187 which is described in Tables 1 and 2 prior to
10 exposure to LPS.

DESCRIPTION OF THE CURRENT EMBODIMENT

This invention is a method for inhibiting cell proliferation disorders, infectious diseases, and immunological diseases in mammals and especially in humans using compositions having the following general formula:




10


In the composition, R_1 - R_4 are each individually selected from the group of compounds including hydrogen, halogen, hydroxyl, thiol, lower alkyl, substituted lower alkyl, alkenyl, alkynyl, alkylalkenyl, alkyl alkynyl, alkoxy, alkylthio, acyl, aryloxy, amino, amido, carboxyl, aryl, substituted aryl, heterocycle, heteroaryl, substituted heterocycle, heteroalkyl, cycloalkyl, substituted cycloalkyl, alkylcycloalkyl, alkylcycloheteroalkyl, nitro, or cyano.

In the composition, R_5 - R_7 are each individually selected from the group of compounds including hydrogen, halogen, hydroxyl, thiol, oxo, lower alkyl, substituted lower alkyl, alkenyl, alkynyl, alkylalkenyl, alkyl alkynyl, alkoxy, alkylthio, acyl, aryloxy, amino, amido, carboxyl, aryl, substituted aryl, heterocycle, heteroaryl, substituted heterocycle, heteroalkyl, cycloalkyl, substituted cycloalkyl, alkylcycloalkyl, alkylcycloheteroalkyl, nitro, or cyano.

X is selected from the group of compounds including hydrogen, $-D_1$, $-D_2$, $-E$, $-D_1-D_2$, $-D_1-E$, $-D_2-E$ or a compound having the formula:




wherein D_1 and D_2 are each individually selected from the group of compounds including



hydrogen, halogen, hydroxyl, thiol, lower alkyl, substituted lower alkyl, alkynyl, alkylalkenyl, alkyl alkynyl, alkoxy, alkylthio, acyl, aryloxy, amino, amido, carboxyl, aryl, substituted aryl, heterocycle, heteroaryl, substituted heterocyclic, heteroalkyl, cycloalkyl, substituted cycloalkyl, alkylcycloalkyl, or alkylcycloheteroalkyl; and

wherein E is selected from the group of compounds including:



hydrogen, halogen, hydroxyl, thiol, lower alkyl, substituted lower alkyl, alkynyl,

alkylalkenyl, alkyl alkynyl, alkoxy, alkylthio, acyl, aryloxy, alkylthio, acyl, aryloxy, amino, amido, carboxyl, aryl, substituted aryl, heterocycle, heteroaryl, substituted heterocycle, heteroalkyl, cycloalkyl, substituted cycloalkyl, alkylcycloalkyl, or alkylcycloheteroalkyl.

- 5 In the compounds identified above, R_5 - R_6 are each individually selected from the group of compounds including hydrogen, halogen, hydroxyl, thiol, oxo, lower alkyl, substituted lower alkyl, alkenyl, alkynyl, alkylalkenyl, alkyl alkynyl, alkoxy, alkylthio, acyl, aryloxy, amino, amido, carboxyl, aryl, substituted aryl, heterocycle, heteroaryl, substituted heterocycle, heteroalkyl, cycloalkyl, substituted cycloalkyl, alkylcycloalkyl,
10 alkylcycloheteroalkyl, nitro, or cyano.

- When D_1 , D_2 and/or E are selected from compounds including substituents R_{10} - R_{14} , J_1 and J_2 , then: R_{10} - R_{14} are each individually selected from the group of compounds including hydrogen, halogen, hydroxyl, thiol, oxo, lower alkyl, substituted lower alkyl, alkenyl, alkynyl, alkylalkenyl, alkyl alkynyl, alkoxy, alkylthio, acyl, aryloxy, amino, amido,
15 carboxyl, aryl, substituted aryl, heterocycle, heteroaryl, substituted heterocycle, heteroalkyl, cycloalkyl, substituted cycloalkyl, alkylcycloalkyl, alkylcycloheteroalkyl, nitro, or cyano; and J_1 and J_2 are each individually selected from the group including N- R_{15} , CR_{16} , R_{17} , O, S-(O)₀₋₂, P-(O)₀₋₃, wherein R_{15} - R_{17} are each individually substituted with a component selected from the group including hydrogen, halogen, hydroxyl, oxo, thiol, lower
20 alkyl, substituted lower alkyl, alkynyl, alkylalkenyl, alkyl alkynyl, alkoxy, alkylthio, acyl, aryloxy, amino, amido, carboxyl, aryl, substituted aryl, heterocyclic, heteroaryl, substituted heterocycle, heteroalkyl, cycloalkyl, substituted cycloalkyl, alkylcycloalkyl, alkylcycloheteroalkyl, or cyano.

The following terms are used to describe various constituents of the chemical composition useful in the method of this invention. The terms are defined as follows:

The term "halogen" refers to fluorine, bromine, chlorine, and iodine atoms.

The term "hydroxyl" refers to the group -OH.

5 The term "oxo" refers to the group =O.

The term "thiol" or "mercapto" refers to the group -SH, and -S(O)₀₋₂.

The term "lower alkyl" refers to a cyclic, branched or straight chain, alkyl group of one to ten carbon atoms. This term is further exemplified by such groups as methyl, ethyl, n-propyl, i-propyl, n-butyl, t-butyl, i-butyl (or 2-methylpropyl), cyclopropylmethyl, i-amyl,
10 n-amyl, hexyl and the like.

The term "substituted lower alkyl" refers to lower alkyl as just described including one or more groups such as hydroxyl, thiol, alkylthiol, halogen, alkoxy, amino, amido, carboxyl, cycloalkyl, substituted cycloalkyl, heterocycle, cycloheteroalkyl, substituted cycloheteroalkyl, acyl, carboxyl, aryl, substituted aryl, aryloxy, hetaryl, substituted hetaryl,
15 aralkyl, heteroaralkyl, alkyl alkenyl, alkyl alkynyl, alkyl cycloalkyl, alkyl cycloheteroalkyl, cyano. These groups may be attached to any carbon atom of the lower alkyl moiety.

The term "alkenyl" refers to a group -CR'=CR''R''' where R', R'', R''' are each selected from hydrogen, halogen, lower alkyl, substituted lower alkyl, acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or the like as defined.

20 The term "alkynyl" refers to a group -C≡C-R'; where R' is selected from hydrogen, halogen, lower alkyl, substituted lower alkyl, acyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl or the like as defined.

The term "alkyl alkenyl" refers to a group -R-CR'=CR''R''', where R is lower alkyl,

or substituted lower alkyl, R', R'', R''' are each independently selected from hydrogen, halogen, lower alkyl, substituted lower alkyl, acyl, aryl, substituted aryl, hetaryl, or substituted hetaryl as defined below.

The term "alkyl alkynyl" refers to a group $-RC\equiv CR'$ where R is lower alkyl or substituted lower alkyl, R' is hydrogen, lower alkyl, substituted lower alkyl, acyl, aryl, substituted aryl, hetaryl, or substituted hetaryl as defined below.

The term "alkoxy" refers to the group $-OR$, where R is lower alkyl, substituted lower alkyl, acyl, aryl, substituted aryl, aralkyl, substituted aralkyl, heteroalkyl, heteroarylalkyl, cycloalkyl, substituted cycloalkyl, cycloheteroalkyl, or substituted cycloheteroalkyl as defined below.

The term "alkylthio" denotes the group $-SR$, $-S(O)_{n-1,2}-R$, where R is lower alkyl, substituted lower alkyl, aryl, substituted aryl aralkyl or substituted aralkyl as defined below.

The term "acyl" refers to groups $-C(O)R$, where R is hydrogen, lower alkyl, substituted lower alkyl, aryl, substituted aryl and the like as defined below.

The term "aryloxy" refers to groups $-OAr$, where Ar is an aryl, substituted aryl, heteroaryl, or substituted heteroaryl group as defined below.

The term "amino" refers to the group NRR' , where R and R' may independently be hydrogen, lower alkyl, substituted lower alkyl, aryl, substituted aryl, hetaryl, cycloalkyl, or substituted hetaryl as defined below or acyl.

The term "amido" refers to the group $-C(O)NRR'$, where R and R' may independently be hydrogen, lower alkyl, substituted lower alkyl, aryl, substituted aryl, hetaryl, substituted hetaryl as defined below.

The term "carboxyl" refers to the group $-C(O)OR$, where R may independently be

hydrogen, lower alkyl, substituted lower alkyl, aryl, substituted aryl, hetaryl, substituted hetaryl and the like as defined.

The terms "aryl" or "Ar" refer to an aromatic carbocyclic group having at least one aromatic ring (e.g., phenyl or biphenyl) or multiple condensed rings in which at least one ring
5 is aromatic, (e.g., 1,2,3,4-tetrahydronaphthyl, naphthyl, anthryl, or phenanthryl).

The term "substituted aryl" refers to aryl optionally substituted with one or more functional groups, e.g., halogen, lower alkyl, lower alkoxy, lower alkylthio, trifluoromethyl, amino, amido, carboxyl, hydroxyl, aryl, aryloxy, heterocycle, hetaryl, substituted hetaryl, nitro, cyano, alkylthio, thiol, sulfamido and the like.

10 The term "heterocycle" refers to a saturated, unsaturated, or aromatic carbocyclic group having a single ring (e.g., morpholino, pyridyl or furyl) or multiple condensed rings (e.g., naphthpyridyl, quinoxalyl, quinolinyl, indolizinyll or benzo[b]thienyl) and having at least one hetero atom, such as N, O or S, within the ring, which can optionally be unsubstituted or substituted with, e.g., halogen, lower alkyl, lower alkoxy, lower alkylthio,
15 trifluoromethyl, amino, amido, carboxyl, hydroxyl, aryl, aryloxy, heterocycle, hetaryl, substituted hetaryl, nitro, cyano, alkylthio, thiol, sulfamido and the like.

The terms "heteroaryl" or "hetar" refer to a heterocycle in which at least one heterocyclic ring is aromatic.

The term "substituted heteroaryl" refers to a heterocycle optionally mono or poly
20 substituted with one or more functional groups, e.g., halogen, lower alkyl, lower alkoxy, lower alkylthio, trifluoromethyl, amino, amido, carboxyl, hydroxyl, aryl, aryloxy, heterocycle, hetaryl, substituted hetaryl, nitro, cyano, alkylthio, thiol, sulfamido and the like.

The term "aralkyl" refers to the group -R-Ar where Ar is an aryl group and R is lower alkyl or substituted lower alkyl group. Aryl groups can optionally be unsubstituted or substituted with, e.g., halogen, lower alkyl, alkoxy, alkyl thio, trifluoromethyl, amino, amido, carboxyl, hydroxyl, aryl, aryloxy, heterocycle, hetaryl, substituted hetaryl, nitro, cyano, alkylthio, thiol, sulfamido and the like.

The term "heteroalkyl" refers to the group -R-Het where Het is a heterocycle group and R is a lower alkyl group. Heteroalkyl groups can optionally be unsubstituted or substituted with e.g., halogen, lower alkyl, lower alkoxy, lower alkylthio, trifluoromethyl, amino, amido, carboxyl, hydroxyl, aryl, aryloxy, heterocycle, hetaryl, substituted hetaryl, nitro, cyano, alkylthio, thiol, sulfamido and the like.

The term "heteroarylalkyl" refers to the group -R-HetAr where HetAr is an heteroaryl group and R lower alkyl or substituted loweralkyl. Heteroarylalkyl groups can optionally be unsubstituted or substituted with, e.g., halogen, lower alkyl, substituted lower alkyl, alkoxy, alkylthio, aryl, aryloxy, heterocycle, hetaryl, substituted hetaryl, nitro, cyano, alkylthio, thiol, sulfamido and the like.

The term "cycloalkyl" refers to a divalent cyclic or polycyclic alkyl group containing 3 to 15 carbon. For polycyclic groups, these may be multiple condensed rings in which one of the distal rings may be aromatic (e.g., indanyl, tetrahydronaphthalene, etc. . . .).

The term "substituted cycloalkyl" refers to a cycloalkyl group comprising one or more substituents with, e.g., halogen, lower alkyl, substituted lower alkyl, alkoxy, alkylthio, aryl, aryloxy, heterocycle, hetaryl, substituted hetaryl, nitro, cyano, alkylthio, thiol, sulfamido and the like.

The term "cycloheteroalkyl" refers to a cycloalkyl group wherein one or more of the

ring carbon atoms is replaced with a heteroatom (e.g., N, O, S or P).

The term "substituted cycloheteroalkyl" refers to a cycloheteroalkyl group as herein defined which contains one or more substituents, such as halogen, lower alkyl, lower alkoxy, lower alkylthio, trifluoromethyl, amino, amido, carboxyl, hydroxyl, aryl, aryloxy, heterocycle, hetaryl, substituted hetaryl, nitro, cyano, alkylthio, thiol, sulfamido and the like.

The term "alkyl cycloalkyl" refers to the group -R-cycloalkyl where cycloalkyl is a cycloalkyl group and R is a lower alkyl or substituted lower alkyl. Cycloalkyl groups can optionally be unsubstituted or substituted with e.g. halogen, lower alkyl, lower alkoxy, lower alkylthio, trifluoromethyl, amino, amido, carboxyl, hydroxyl, aryl, aryloxy, heterocycle, hetaryl, substituted hetaryl, nitro, cyano, alkylthio, thiol, sulfamido and the like.

The term "amino acid" refers to the D- or L- isomer of naturally occurring and synthetic alpha amino acids preferably the amino acids are naturally occurring alanine, arginine, asparagine, aspartic acid, cysteine, glutamine, glutamic acid, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine or valine.

Typically D₁, D₂ and E, if present in the composition, may be amino acids. Usually lipophilic amino acids are preferred. In general, the amino acid abbreviations follow the IUPAC-IUB Joint Commission on Biochemical Nomenclature as described in Eur. J. Biochem, 158, 9 (1984).

It is preferred that R₁ is methoxy, -D₁ is leucine, and -D₂ is leucine, and E is NR'R''. It is most preferred that -D₁ is l-leucine, and -D₂ is d-leucine and that E is selected from the group of compounds consisting of benzylamine, 1-indanylamine, N,N'-dibenzylamine, 2,6-difluorobenzylamine, 4-methoxybenzylamine, piperonyl amine, NH₂ and glycineamide.

In one preferred composition, R₃ is methoxy, D₁ is leucine, D₂ is leucine, and E is benzylamine. In another preferred composition, R₃ is methoxy, D₁ is leucine, D₂ is leucine, and E is 1-indanylamine. In still another preferred composition, R₃ is methoxy, D₁ is leucine, D₂ is leucine, and E is N,N-dibenzylamine. In another preferred composition, R₃ is methoxy, D₁ is leucine, D₂ is leucine, and E is 2,6-difluorobenzylamine.

In these preferred compositions, it is further preferred that D₁ is L-Leucine and D₂ is D-Leucine. Known compounds that may be useful in the therapeutic method of this invention are disclosed in Table 1 immediately below.

Table 1

R1	R2	R3	R4	R5	R6	R7	R8	R9	D1
	Cl			Me					OH
		OMe	OMe				=N-OH		OMe
				Ph					Me
	Cl			Me			=N-OH		OEt
	OMe								OH
							Me		OMe
F		F							NH2
									NHEt
					Ph		=CHPh		OH
		F							OH
									OH
		OMe		Me					OH
	Me			Me					OH
				Ph			=N-OH		OMe
		OMe					=CH-(4-dimethylaminophenyl)		OH

R1	R2	R3	R4	R5	R6	R7	R8	R9	D1
	OMe						=N-OH		OMe
				Ph			COOEt		Me
				Me				=N-OH	OEt
		Me							OMe
					Me				OH
	Cl			Me			NHCOCOO Et		OEt
		OMe	OMe						OH
				Me					Me
F		F							OH
									NHMe
				Ph					OH
		OMe		Me				=N-OH	OMe
		OMe							OH
				Me			NHAc		OMe
					Ph		Ph		OH
				Ph					NHCyclopropyl
	OMe	OMe							OH
				Ph					OH
				COO Me					OMe
	OMe	OMe						=N-OH	OMe
				Ph					OMe
	Cl			Me					OEt
		OMe	OM E						OMe
							Me		OH

R1	R2	R3	R4	R5	R6	R7	R8	R9	D1
				Me			NHCOCOO Et		OEt
									NH-n-propyl
OMe			NO2						OH
		F							Cl
		OMe							OMe
		OMe		Me			=N-OH		OMe
		OMe		Me					OMe
	Me			Me					OMe
		F					Me		OH
					4-Methoxyphenyl				OH
							=N-OH		OMe
				Ph	CN		COOEt		OEt
				Me					OH
	OMe								OMe
		OMe							4-[[4-(1-methylethyl)-1-piperazinyl]carbonyl]phenoxy
	Cl			Me			NH2		OEt
		Me							OH
							Me		OH
F		F							Cl
									OMe

R1	R2	R3	R4	R5	R6	R7	R8	R9	D1
				Ph					OH
		OMe						=N-OH	OMe
					2-naphthyl				OH
		OMe		Me					OMe
					Ph		Ph		OMe
				Ph					NH-n-propyl
		Me						=N-OH	OMe
					Ph				OH
		OMe						=N-OH	OMe
				Ph	SH		COOEt		OH
	Cl			Me					OH
	OMe	OMe							OMe
					Me				OH
				Me			NH2		OEt
									NH-n-Butyl
	NO2	OMe							OH
		F							NH2
							Me		OH
				Ph					OH
		OMe							OMe
	Me			Me			3-propanoic acid	3-propanoic acid	OH
				Me					OMe
	OMe				Ph				OH
		OMe		Ph					NHMe

R1	R2	R3	R4	R5	R6	R7	R8	R9	D1
				Me			=N-OH		OMe
				Ph	COO Et		COOEt		OEt
		OMe		COO Me					OMe

It is within the knowledge of one of skill in the art that stereoisomers of the compositions described herein as well as isomer and stereoisomers of components that comprise the compositions identified herein all fall within the scope of compositions that are
5 useful in the therapeutic method of this invention.

If the compound useful in the method of this invention contains a basic group, an acid addition salt may be prepared. Acid addition salts of the compounds are prepared in a standard manner in a suitable solvent from the parent compound and an excess of acid, such as hydrochloric, hydrobromic, sulfuric, phosphoric, acetic, maleic, succinic, or
10 methanesulfonic. If the final compound contains an acidic group, cationic salts may be prepared. Typically the parent compound is treated with an excess of an alkaline reagent, such as hydroxide, carbonate or alkoxide, containing the appropriate cation. Cations such as Na^+ , K^+ , Ca^{+2} and NH_4^{+} are examples of cations present in pharmaceutically acceptable salts. Certain of the compounds form inner salts or zwitterions which may also be acceptable.

15 The compounds described above are useful for treating cell proliferation disorders, infectious diseases, and immunological diseases in mammals, and specifically, in human patients who require such treatment. Cell proliferative disorders that may be treated using the composition described above include cancer, cardiovascular disease such as myocarditis and restenosis following angioplasty, renal diseases such as lupus and polycystic kidney disease.

host graft rejection, gout, and other proliferative disorders. Autoimmune diseases that may be treated with the compositions described above include rheumatoid arthritis, lupus, type I diabetes, multiple sclerosis and similar disorders and diseases. Infectious diseases that may be treated using the compositions described above include IBD, Crohn's disease, AIDS, ARDS, and similar disorders. The compositions disclosed above can also be used to treat fungal infections, dermatological diseases such as psoriasis, abnormal wound healing, keloids, immunological diseases such as autoimmunity, asthma, allergy, acute and delayed hypersensitivity, graft versus host disease, transplant rejection, and neuroimmunological disease such as multiple sclerosis and acute disseminated encephalomyelitis.

10 The method of treatment of these diseases and disorders comprises the administration parenterally, or orally of an effective quantity of the chosen compound or combinations thereof, preferably dispersed in a pharmaceutical carrier. Dosage units of the active ingredient are generally selected from the range of 0.01 to 100 mg/kg, but will be readily determined by one skilled in the art depending upon the route of administration, age and
15 condition of the patient. These dosage units may be administered one to ten times daily for acute or chronic disease. No unacceptable toxicological effects are expected when compounds of the invention are administered in accordance with the present invention.

Pharmaceutical compositions of the compounds of this invention, or derivatives thereof, may be formulated as solutions or lyophilized powders for parenteral
20 administration. Powders may be reconstituted by addition of a suitable diluent or other pharmaceutically acceptable carrier prior to use. The liquid formulation is generally a buffered, isotonic, aqueous solution. Examples of suitable diluents are normal isotonic saline solution, standard 5% dextrose in water or buffered sodium or ammonium acetate

solution. Such formulation is especially suitable for parenteral administration, but may also be used for oral administration. It may be desirable to add excipients such as polyvinylpyrrolidinone, gelatin, hydroxycellulose, acacia, polyethylene glycol, mannitol, sodium chloride or sodium citrate. Alternatively, these compounds may be encapsulated, tableted or prepared in an emulsion or syrup for oral administration. Pharmaceutically acceptable solid or liquid carriers may be added to enhance or stabilize the composition, or to facilitate preparation of the composition. Liquid carriers include syrup, peanut oil, olive oil, glycerin, saline, alcohols and water. Solid carriers include starch, lactose, calcium sulfate, dehydrate, terra alba, magnesium stearate or stearic acid, talc, pectin, acacia, agar or gelatin. The carrier may also include a sustained release material such as glyceryl monostearate or glyceryl distearate, alone or with a wax. The amount of solid carrier varies but, preferably, will be between about 20 mg to about 1 g per dosage unit. The pharmaceutical preparations are made following the conventional techniques of pharmacy involving milling, mixing, granulation, and compressing, when necessary, for tablet forms; or milling, mixing and filling for hard gelatin capsule forms. When a liquid carrier is use, the preparation will be in the form of a syrup, elixir, emulsion or an aqueous or non-aqueous suspension. Such a liquid formulation may be administered directly p.o. or filled into a soft gelatin capsule.

EXAMPLE 1

The compounds useful in the therapeutic method of this invention are prepared by conventional methods of organic chemistry. Coupling reagents are well known in the art, such as DCC and other carbodiimides, EDC, BOP and PPA, and they may optionally
5 be used with other reagents, such as HOBT, NMM and DMAP, which may facilitate the reaction. Preparation of compounds of the formula (1) wherein D₁, D₂ and E are amino acids are well known in the art using either conventional solution phase or solid phase techniques as described in Bodanszky, "The Practice of Peptide Synthesis," Springer-Verlag, First Edition, 1984. Suitable protective groups for the amino group are
10 those disclosed by Greene et al., "Protective Group in Organic Synthesis," Second Edition, John Wiley and Sons, New York, 1991. The benzyloxycarbonyl, t-butoxycarbonyl and fluorenylmethoxycarbonyl groups are especially useful amino protecting groups.

Solid phase peptide synthesis was accomplished as follows. Rink amide resin is placed in a syringe fitted with a fritted filter. The resin is deprotected using 20% piperidine
15 in DMF. After 20 minutes, the resin was washed five times with DMF, five times with methanol, then five times with DMF. A solution of amino acid (E), carbodiimide and HOBT in DMF was drawn up into the syringe and the reaction mixture was allowed to mix for 4 to 20 hours. The reaction solution was ejected and the mixture washed five times with DMF, five times with methanol, then five times with DMF. This cycle was
20 repeated until the desired sequence was attached. The final coupling used, 5-methoxy-1-indanone-3-acetic acid, carbodiimide and HOBT. After final washings of the resin, the peptide fragment was cleaved from the resin using 95% TFA/5% water. Concentration of the cleavage mixture affords a white solid.

EXAMPLE 2

Compounds of this invention prepared according to the method of Example 1 were tested as follows. The 20S catalytic subunit of the proteasome (also known as the multicatalytic proteinase complex) was purified to homogeneity from bovine brain according to published methods (Wilk S. and Orlowski, M, 40 842 J. Neurochem (1983)). The chymotryptic activity of the complex is measured by the increase in fluorescence following cleavage of the substrate peptide succinyl-leucine-leucine-valine-tyrosine-7-amino-4-methyl coumarin. The standard *in vitro* assay consists of 2µg 20S proteasome, 0.1 - 100µg/ml proteasome inhibitor in 200µl 50mM HEPES, containing 0.1 % sodium dodecyl sulphate, pH7.5. The proteolytic reaction is initiated by the addition of 50µM fluorogenic peptide substrate and allowed to progress for 15 minutes at 37°C. The reaction is terminated by the addition of 100µl of 100mM acetate buffer, pH4.0. The rate of proteolysis is directly proportional to the amount of liberated aminomethylcoumarin which is measured by fluorescent spectroscopy (EX 370nm, EM 430nm). The structures of the compounds tested as well as the test results are reported in Table 2, below.

TABLE 2

Compound	R3	D1	D2	E	IC50 ug/ml
1	MeO	d-leu-NH2			5
2	MeO	d-leu	d-leu-NH2		1
3	MeO	leu-NH2			>10
4	MeO	d-leu	his-NH2		>10
5	MeO	leu	leu-NH2		>10
6	MeO	d-leu	leu-NH2		>10

Compound	R3	D1	D2	E	IC50 ug/ml
7	MeO	leu	d-leu-NH2		>10
8	MeO	Nle-NH2			>10
9	MeO	d-Nle-NH2			>10
10	MeO	Nva-NH2			>10
11	MeO	d-leu	Aib-NH2		>10
12	MeO	d-phe-NH2			>10
13	MeO	CHA	Leu-NH2		>10
14	MeO	phe	leu-NH2		>10
15	MeO	Nle	d-Leu-NH2		>10
16	MeO	Nle	leu-NH2		>10
17	MeO	leu	leu	gly-NH2	>10
18	MeO	d-leu	leu	gly-NH2	5
19	MeO	d-leu	d-leu	gly-NH2	>10
20					
21	MeO	leu	d-leu	gly-NH2	0.1
22	MeO	d-Nle	d-leu-NH2		>10
23	MeO	d-Nle	leu-NH2		>10
24	MeO	phe-NH2			>10
25	MeO	Tic-NH2			>10
26	MeO	Tic	d-leu-NH2		>10
27	MeO	d-phe	d-leu-NH2		10
28	MeO	leu	Aib-NH2		>10
29	MeO	CHA-NH2			>10
30	MeO	d-val	d-leu-NH2		>10
31	MeO	d-pro	d-leu-NH2		10

Compound	R3	D1	D2	E	IC50 ug/ml
32	MeO	d-cha	d-leu-NH2		>10
33	MeO	d-leu	d-val-NH2		>10
34	MeO	d-val	d-val-NH2		>10
35	MeO	d-cha	d-cha-NH2		>10
36	MeO	d-phg-NH2			>10
37	MeO	d-phg	d-leu-NH2		>10
38	MeO	benzyl			>10
39	MeO	dibenzylamin e			>10
40	MeO	gly	gly	gly-NH2	>10
41	MeO	leu	gly	gly-NH2	>10
42	MeO	gly	d-leu	gly-NH2	>10
43	MeO	nle	d-leu	gly-NH2	>10
44	MeO	nva	d-leu	gly-NH2	>10
45	MeO	phe	d-leu	gly-NH2	>10
46	MeO	cha	d-leu	gly-NH2	>10
47	MeO	val	d-leu	gly-NH2	>10
48	MeO	phg	d-leu	gly-NH2	>10
49	MeO	pro	d-leu	gly-NH2	>10
50	MeO	leu	d-leu	gly-NH2	0.5
51	MeO	leu	d-leu	gly-NH2	0.5
52	MeO	d-leu	ala-NH2		>10
53	MeO	d-leu	d-ala-NH2		>10
54	MeO	d-leu	d-pro-NH2		>10
55	MeO	d-pro-NH2			>10
56	MeO	d-leu	d-phe-NH2		>10

Compound	R3	D1	D2	E	IC50 ug/ml
57	MeO	d-nva	d-leu-NH2		>10
58	MeO	nva	d-leu-NH2		>10
59	MeO	d-ala	d-leu-NH2		>10
60	MeO	d-tic	d-leu-NH2		>10
61	MeO	d-ser	d-leu-NH2		>10
62	MeO	diisopropyl			>10
63	MeO	morpholine			>10
64	MeO	pyrrolidine			>10
65	MeO	phenethylamine			>10
66	MeO	phenpropylamine			>10
67	MeO	piperidine			>10
68	MeO	diethylamine			>10
69	MeO	cyclobutylamine			>10
70	MeO	heptylamine			>10
71	MeO	3-methoxypropylamine			>10
72	MeO	3,4-dimethoxyphenethylamine			>10
73	MeO	N-methylbenzylamine			>10
74	MeO	cyclopentylamine			>10
75	MeO	2,6-dimethylpiperidine			>10

Compound	R3	D1	D2	E	IC50 ug/ml
76	MeO	N-benzylethanol amine			>10
77	MeO	indoline			>10
78	MeO	dimethylamin e			>10
79	MeO	bis(2-methoxyethyl) amine			>10
80	MeO	piperonylami ne			>10
81	MeO	4-hydroxypiperi dine			>10
82	MeO	3-iodoaniline			>10
83	MeO	1-aminoindane			>10
84	MeO	ethanolamine			>10
85	MeO	4-methoxybenz ylamine			>10
86	MeO	leu	d-nle	gly-NH2	>10
87	MeO	leu	d-nva	gly-NH2	>10
88	MeO	leu	d-phe	gly-NH2	>10
89	MeO	leu	d-cha	gly-NH2	>10
90	MeO	leu	d-val	gly-NH2	>10
91	MeO	leu	d-phg	gly-NH2	10
92	MeO	leu	d-pro	gly-NH2	>10
93	MeO	d-arg	d-leu-NH2		>10
94	MeO	d-asp	d-leu-NH2		>10

Compound	R3	D1	D2	E	IC50 ug/ml
95	MeO	d-asn	d-leu-NH2		10
96	MeO	d-asn	d-leu-NH2		>10
97	MeO	d-glu	d-leu-NH2		>10
98	MeO	d-gln	d-leu-NH2		>10
99	MeO	d-his	d-leu-NH2		>10
100	MeO	d-lys	d-leu-NH2		>10
101	MeO	d-thi	d-leu-NH2		>10
102	MeO	d-tyr	d-leu-NH2		>10
103	MeO	d-trp	d-leu-NH2		>10
104	MeO	4- (aminomethyl) pyridine			>10
105	MeO	1,2- diaminopropane			>10
106	MeO	thiomorpholine			>10
107	MeO	2- methoxybenzylamine			>10
108	MeO	4- methylpiperidine			>10
109	MeO	3-pyrrolidinol			>10
110	MeO	4-amino-1- benzylpiperidine			>10
111	MeO	3-amino-1,2- propandiol			>10
112	MeO	1-(2- aminoethyl)p			>10

Compound	R3	D1	D2	E	IC50 ug/ml
		pyrrolidine			
113	MeO	2-amino-2-methyl-1-propanol			>10
114	MeO	2-(amionmethyl)pyridine			>10
115	MeO	2-(methy lamio) ethanol			>10
116	MeO	3-(3-pyridylmethy lamio)propio nitrile			>10
117	MeO	2-methyoxethyl amine			>10
118	MeO	2-amino-1-phenyl-1,3-propanediol			>10
119	MeO	2-pyrrolidinom ethanol			>10
120	MeO	3-phenyl-1-propylamine			>10
121	MeO	p-anisidine			>10
122	MeO	aniline			>10
123	MeO	leu	d-leu	d-val-NH2	>10
124	MeO	leu	d-leu	val-NH2	>10
125	MeO	leu	d-leu	d-ala-NH2	>10
126	MeO	leu	d-leu	ala-NH2	>10
127	MeO	leu	d-leu	d-phe-NH2	>10

Compound	R3	D1	D2	E	IC50 ug/ml
128	MeO	leu	d-leu	phe-NH2	>10
129	MeO	d-leu	d-arg-NH2		>10
130	MeO	d-leu	d-asp-NH2		>10
131	MeO	d-leu	d-asn-NH2		>10
132	MeO	d-leu	d-cha-NH2		>10
133	MeO	d-leu	d-glu-NH2		>10
134	MeO	d-leu	d-nle-NH2		>10
135	MeO	d-leu	d-tyr-NH2		>10
136	MeO	d-leu	d-trp-NH2		>10
137	MeO	d-leu	d-gln-NH2		>10
138	MeO	d-leu	d-lys-NH2		>10
139	MeO	d-leu	d-nva-NH2		>10
140	MeO	d-leu	d-phg-NH2		>10
141	MeO	d-leu	d-ser-NH2		>10
142	MeO	d-leu	d-thi-NH2		>10
143	MeO	d-leu	d-tic-NH2		>10
144	MeO	N-(4-hydroxyphenyl)-2-naphthylamine			>10
145	MeO	2-amino-4,6-dihydroxy-5-methylpyrimidine			>10
146	MeO	5-(hydroxymethyl)-2-pyrrolidinone			>10
147	MeO	3-hydroxydiphe			>10

Compound	R3	D1	D2	E	IC50 ug/ml
		nylamine			
148	MeO	2-amino-4-phenylphenol			>10
149	MeO	2-(4-methoxybenzylamino)pyridine			>10
150	MeO	hexamethylenediamine			>10
151	MeO	4-hydroxy-4-phenylpiperidine			>10
152	MeO	4-iodoaniline			>10
153	MeO	2-methyl-6-nitroaniline			>10
154	MeO	(r)-(-)-5-(hydroxymethyl)-2-pyrrolidinone			>10
155	MeO	2-amino-4-chloro-6-methylpyrimidine			>10
156	MeO	2-amino-5-chloropyridine			>10
157	MeO	3,4-dichloroaniline			>10
158	MeO	4-amino-2-mercaptopyrimidine			>10
159	H	leu	d-leu	gly-NH2	>10
160	MeO	indoline			>10

Compound	R3	D1	D2	E	IC50 ug/ml
161	MeO	phenpropyl			>50
162	MeO	p-anisidine			>50
163	MeO	piperonyl			>50
164	MeO	2-pyrrolidine- methanol			>50
165	MeO	2-amino-1- phenyl-1,3- propanediol			>50
166	MeO	N-benzyl- ethanolamine			>50
167	MeO	dimethylamid e			>50
168	MeO	anilino			>50
169	MeO	bis-2- methoxyethyl amine			>50
170	MeO	leu	d-leu	aib-NH2	>10
171	MeO	leu	d-leu-OH		>10
172	MeO	leu	d-leu-OH		>10
173	MeO	leu	d-leu	benzyl	0.5
174	MeO	leu	d-leu	morpholine	>10
175	MeO	leu	d-leu	piperdiny	>10
176	MeO	leu	d-leu	pyrrolidino	>10
177	MeO	leu	d-leu	dibenzyl	1
178	MeO	leu	d-leu	hydroxyethyl	>10
179	MeO	leu	d-leu	N-methylbenzyl	>10
180	MeO	leu	d-leu	N-methylbenzyl	>10
181	MeO	aminomethyl cyclohexanea			>10

Compound	R3	D1	D2	E	IC50 ug/ml
		mide			
182	MeO	leu	aminomethylcyclohexaneamide		>10
183	MeO	leu	aminomethylcyclohexaneamide	gly-NH2	>10
184	MeO	leu	d-leu	phenyl	>10
185	MeO	leu	d-leu	phenethyl	>10
186	MeO	leu	d-leu	phenpropyl	>10
187	MeO	leu	d-leu	indane	0.5
188	MeO	leu	d-leu	aminomethylcyclohexane	>10
189	MeO	leu	d-leu	aminomethylpyridine	>10
190	MeO	leu	d-leu	adamantyl	>10
191	MeO	leu	d-leu	tetrahydroisoquinoline	>10
192	MeO	leu	d-leu	4-pyridylmethyl	10
193	MeO	leu	d-leu	N-benzylhydroxamic	0.4
194	MeO	leu	d-leu	4-methoxybenzyl	>10
195	MeO	leu	d-leu	4-nitrobenzyl	10
196	MeO	leu	d-leu	2,6-difluorobenzyl	0.08
197	MeO	leu	d-leu	piperonyl	0.2
198	MeO	d-leu	benzyl		>10
199	MeO	d-leu	dibenzyl		>10
200	MeO	d-leu	isoamyl		>10

EXAMPLE 3

Compounds prepared according to the method of Example 1 were assayed against several different cell lines. Cell monolayers were cultured in the presence of test compound
5 for 18 hours to assess their ability to inhibit cell proliferation. Cell proliferation was determined colorimetrically using the Celltiter 96 Aqueous non-radioactive cell proliferation assay (Promega) where cell proliferation is directly proportional to absorbance at 490nm. Results are quoted as the IC₅₀ in µg/ml for inhibition of cell proliferation in various cell types.

Compound	RAW	MCF-7	OVCAR	CaCo	Panc-I
173	20	10	8	17	17
187	10	6	5	8	7
194	1		3		8
196	7		2		8
197	12		2		8

EXAMPLE 4

Compounds prepared according to the method of Example 1 were tested for the inhibition of LPS induced TNF synthesis. RAW cells were pretreated with different concentrations of test compound for 1 hour prior to the administration of lipopolysaccharide (100ng/ml). Cell culture supernatants were harvested after 1 hour and assayed for TNF concentration by ELISA (Biosource)

Compound	IC ₅₀ (µg/ml)
173	5
187	5
194	3
196	3
197	3

EXAMPLE 5

This Example examines the ability of compound 173 and particularly compound 187 described in Tables 1 & 2 above to inhibit proteasome activity as indicated, in part, by the presence of I κ B and/or p105 in inhibited cells. In order for NF- κ B to translocate to the nucleus in response to a stimulus such as lipopolysaccharide (LPS) and activate transcription two proteolytic events need to occur, namely degradation of the inhibitory protein I κ B and processing of p105 to p50. These proteolytic events serve to unmask the nuclear localization signal of NF- κ B.

Inhibition of LPS-induced I κ B Degradation

10 RAW cells were pretreated with different concentrations of test compound for 1 hour prior to the administration of lipopolysaccharide (100ng/ml). Whole cell lysates were harvested after 1 hour, 10 μ g protein was separated by SDS-PAGE, transferred to nitrocellulose and assayed for immunoreactivity with anti-I κ B α antibody. Western blots, (See Figure 1), were visualized using the Boehringer Mannheim Chemiluminescent detection
15 kit. The blot shows that I κ B is present in cells treated with as little as 5 μ g/ml of compounds 173 and 187.

Inhibition of LPS-induced p150 to p50 Processing

Compound 187 as described in Tables 1 & 2 above was used to pretreat RAW cells as described above, and whole cell lysates prepared as described above were analyzed for
20 immunoreactivity toward an anti-p50 antibody. The results, set forth in Figure 2 indicate that p50 and p150 are both present in cells treated with as little as 5 μ g/ml of compound 187, while in untreated cells, majority of P105 has been processed to p50.

Inhibition of LPS-induced Translocation of NF- κ B to the Nuclear Fraction of the


Cell

RAW cells were treated for 1 hour with compound 187 (20 μ g/ml) and then incubated with lipopolysaccharide (100 ng/ml) for a further one hour. Nuclear fractions were prepared according to standard procedures. Binding reactions for gel mobility shift assays contained

5 5 μ g of nuclear extract protein, 50,000 cpm of 32 P-labeled NF- κ B consensus binding oligonucleotide in the presence and absence of a fifty fold excess of unlabeled oligonucleotide. The gel mobility shift assay, set forth in Figure 3 shows that compound 187 is effective in inhibiting the accumulation of NF- κ B in the nucleus of cells.

What we claim is:


1. A method for treating mammals with cell proliferation disorders, immune system disorders, and infectious diseases comprising administering to the mammal a therapeutically effective amount of a compound having the formula:



R_1 - R_4 are each individually selected from the group of compounds including hydrogen, halogen, hydroxyl, thiol, lower alkyl, substituted lower alkyl, alkenyl, alkynyl, alkylalkenyl, alkyl alkynyl, alkoxy, alkylthio, acyl, aryloxy, amino, amido, carboxyl, aryl, substituted aryl, heterocycle, heteroaryl, substituted heterocycle, heteroalkyl, cycloalkyl, substituted cycloalkyl, alkylcycloalkyl, alkylcycloheteroalkyl, nitro, or cyano.


R_5 - R_9 are each individually selected from the group of compounds including hydrogen, halogen, hydroxyl, thiol, oxo, lower alkyl, substituted lower alkyl, alkenyl, alkynyl, alkylalkenyl, alkyl alkynyl, alkoxy, alkylthio, acyl, aryloxy, amino, amido, carboxyl, aryl, substituted aryl, heterocycle, heteroaryl, substituted heterocycle, heteroalkyl, cycloalkyl, substituted cycloalkyl, alkylcycloalkyl, alkylcycloheteroalkyl, nitro, or cyano;

X is selected from the group of compounds including hydrogen, $-D_1$, $-D_2$, $-E$, $-D_1-D_2$, $-D_1-E_1$, $-D_2-E$ or a compound having the formula:



5

wherein D_1 and D_2 are each selected from the group of compounds including:




10

hydrogen, halogen, hydroxyl, thiol, lower alkyl, substituted lower alkyl, alkynyl, alkylalkenyl, alkyl alkynyl, alkoxy, alkylthio, acyl, aryloxy, amino, amido, carboxyl, aryl, substituted aryl, heterocycle, heteroaryl, substituted heterocyclic, heteroalkyl, cycloalkyl, substituted cycloalkyl, alkylcycloalkyl, alkylcycloheteroalkyl, or is absent;

15

wherein E is a compound selected from the group including:



20

hydrogen, halogen, hydroxyl, thiol, lower alkyl, substituted lower alkyl, alkynyl, alkylalkenyl, alkyl alkynyl, alkoxy, alkylthio, acyl, aryloxy, alkylthio, acyl, aryloxy, amino, amido, carboxyl, aryl, substituted aryl, heterocycle, heteroaryl, substituted heterocycle, heteroalkyl, cycloalkyl, substituted cycloalkyl, alkylcycloalkyl, or

alkylcycloheteroalkyl; wherein

R_{10} - R_{14} are each individually selected from the group of compounds including hydrogen, halogen, hydroxyl, thiol, oxo, lower alkyl, substituted lower alkyl, alkenyl, alkynyl, alkylalkenyl, alkyl alkynyl, alkoxy, alkylthio, acyl, aryloxy, amino, amido, carboxyl, aryl, substituted aryl, heterocycle, heteroaryl, substituted heterocycle, heteroalkyl, cycloalkyl, substituted cycloalkyl, alkylcycloalkyl, alkylcycloheteroalkyl, nitro, or cyano;


J_1 and J_2 are $N-R_{15}$, CR_{16} , R_{17} , O, $S-(O)_{0-2}$, $P-(O)_{0-3}$; and

R_{15} - R_{17} are each individually selected from the group of compounds including hydrogen, halogen, hydroxyl, oxo, thiol, lower alkyl, substituted lower alkyl, alkynyl, alkylalkenyl, alkyl alkynyl, alkoxy, alkylthio, acyl, aryloxy, amino, amido, carboxyl, aryl, substituted aryl, heterocyclic, heteroaryl, substituted heterocycle, heteroalkyl, cycloalkyl, substituted cycloalkyl, alkylcycloalkyl, alkylcycloheteroalkyl, or cyano.

2. The method according to claim 1 wherein R_1 - R_4 are each individually selected from the group consisting of hydrogen, halogen, lower alkyl, substituted lower alkyl, aryl, aryloxy, substituted aryl, amino, amido, alkoxy, thio, alkyl thiol, hydroxyl, cyano, nitro, acyl, carboxyl, and alkynyl, and R_5 - R_6 are each individually selected from the group consisting of hydrogen, lower alkyl, substituted lower alkyl, aryl, substituted aryl, amino, amido, alkoxy, and alkynyl.

3. The method according to claim 1 wherein X is selected from the group consisting of $-D_1-E$, or $-D_2-E$.

4. The method according to claim 3 wherein $-D_1$ and $-D_2$ are each:



5. The method according to claim 4 wherein J_1 is $N-R_{15}$, and R_5-R_{11} and R_{15} are each individually selected from the group of compounds consisting of hydrogen, lower alkyl, substituted lower alkyl, aryl, and substituted aryl, and E is selected from the group consisting of lower alkyl, substituted lower alkyl, aryl, substituted aryl, alkoxy, and amino.

6. The method according to claim 5 wherein R_5-R_{11} and R_{15} are each individually selected from the group consisting of hydrogen, lower alkyl, and substituted lower alkyl, and E is selected from the group consisting of lower alkyl, substituted lower alkyl, aryl, substituted aryl, alkoxy, and amino.


7. The method according to claim 6 wherein E is selected from the group consisting of alkoxy and amino, and R_1-R_4 is selected from the group consisting of hydrogen, halogen, lower alkyl, substituted lower alkyl, alkoxy, amino, nitro, hydroxyl, cyano, alkynyl, thio, and alkylthio.

8. The method according to claim 7 wherein E is $NR'R''$ wherein R' and R'' are each individually selected from the group consisting of hydrogen, lower alkyl, substituted lower alkyl, aryl, substituted aryl, and cycloalkyl, and R_1-R_4 are each individually selected from the group consisting of hydrogen, halogen, nitro, lower alkyl, substituted lower alkyl, amino, alkylthio, thio, hydroxy, and alkoxy.


9. The method according to claim 8 wherein R_1-R_4 are each individually selected from the group consisting of hydrogen, alkoxy, and lower alkyl, R_5-R_{11} and R_{15} are each individually selected from the group consisting of hydrogen, lower alkyl, and substituted lower alkyl, and R' and R'' are each individually selected from the group consisting of

hydrogen, lower alkyl, substituted lower alkyl, and cycloalkyl.

10. The method according to claim 2 wherein X is:




11. The method according to claim 10 wherein -D₁ and -D₂ are each:



12. The method according to claim 11 wherein E is a compound selected from the group consisting of hydrogen, lower alkyl, substituted lower alkyl, aryl, substituted aryl, amino, cycloalkyl, substituted cycloalkyl, alkoxy, or alkylcycloalkyl.

13. The method according to claim 11 wherein E is:



14. The method according to claim 13 wherein J_1 and J_2 are each $N R_{15}$, wherein R_{10} - R_{13} and R_{15} are each individually selected from the group consisting of hydrogen, lower alkyl, substituted lower alkyl, aryl, and substituted aryl, and R_{14} is selected from the group consisting of lower alkyl, substituted lower alkyl, aryl, substituted aryl, amino, and alkoxy.

5 15. The method according to claim 14 wherein R_5 - R_{13} are each individually selected from the group consisting of hydrogen, lower alkyl, substituted lower alkyl, aryl, and substituted aryl, R_{14} is selected from the group of compounds consisting of alkoxy, and amino, and R_{15} is selected from the group of compounds consisting of hydrogen, lower alkyl, substituted lower alkyl, aryl, and substituted aryl.

10 16. The method according to claim 15 wherein R_1 - R_4 are each individually selected from the group of compounds consisting of hydrogen, halogen, lower alkyl, substituted lower alkyl, aryl, aryloxy, substituted aryl, amino, alkylthio, nitro, hydroxy, thio, and alkoxy.

15 17. The method according to claim 16 wherein R_5 - R_{13} and R_{15} are each individually selected from the group consisting of hydrogen, lower alkyl, and substituted lower alkyl.

18. The method according to claim 17 wherein R_{14} is $NR'R''$ where R' and R'' are each individually selected from the group of compounds consisting of hydrogen, lower alkyl, substituted lower alkyl, aryl, substituted aryl, and cycloalkyl.

20 19. The method according to claim 18 wherein R_1 - R_4 are each individually selected from the group consisting of hydrogen, alkoxy, and lower alkyl, R_5 - R_{13} and R_{15} are each individually selected from the group of compounds consisting of hydrogen, lower alkyl, and substituted lower alkyl, and R' and R'' are each individually selected from the group of compounds consisting of hydrogen, lower alkyl, cycloalkyl, and substituted lower alkyl.

20. The method according to claim 12 wherein J_1 is $-N R_{15}$, wherein R_{15} is selected from the group of compounds consisting of hydrogen, lower alkyl, substituted lower alkyl, aryl, and substituted aryl.

21. The method according to claim 20 wherein R_5 - R_{11} are each individually
5 selected from the group consisting of hydrogen, lower alkyl, substituted lower alkyl, aryl, and substituted aryl; E is selected from alkoxy and amino, R_1 - R_4 are each individually selected from the group of compounds consisting of hydrogen, halogen, lower alkyl, substituted lower alkyl, alkyl thio, thio, alkoxy, amino, nitro, and hydroxyl, and R_{15} is a compound selected from the group of compounds consisting of hydrogen, lower alkyl, and aryl.

10 22. The method according to claim 21 wherein E is $NR'R''$ wherein R' and R'' are each individually selected from the group of compounds consisting of hydrogen, lower alkyl, substituted lower alkyl, aryl, substituted aryl, and cycloalkyl.

23. The method according to claim 21 wherein R_1 - R_4 are each individually
selected from the group consisting of hydrogen, alkoxy, and lower alkyl, R_5 - R_{11} are each
15 individually selected from the group consisting of hydrogen, lower alkyl, and substituted lower alkyl, R' and R'' are each individually selected from the group consisting of hydrogen lower alkyl, substituted lower alkyl, and cycloalkyl, and R_{15} is selected from the group of compounds selected from the group consisting of hydrogen and lower alkyl.

24. The method according to claim 10 wherein R_3 is methoxy, $-D_1$ is leucine, and
20 $-D_2$ is leucine, and E is $NR'R''$.

25. The method according to claim 24 wherein $-D_1$ is l-leucine, and $-D_2$ is d-leucine.

26. The method according to claim 25 wherein E is selected from the group of compounds consisting of benzylamine, 1-indanylamine, N,N'-dibenzylamine,

2,6-difluorobenzylamine, 4-methoxybenzylamine, piperonyl amine, and NH_3 .

27. The method according to claim 24 wherein E is glycineamide.

28. The method according to claim 1 wherein the mammal is a human.

29. The method according to claim 1 wherein the therapeutically effective amount
5 ranges from about 0.001 to about 100 mg/kg weight of the mammal.

30. The method of claim 1 wherein the composition is administered to a mammal
suffering from a cell proliferation disorder selected from the group consisting of
rheumatoid arthritis, lupus, type I diabetes, multiple sclerosis, cancer, restenosis, host graft
disease, and gout.

10 31. The method of claim 30 wherein the cell proliferation disorder is restenosis.

32. The method of claim 30 wherein the cell proliferation is disorder cancer.

33. The method according to claim 30 wherein the therapeutic agent induces
apoptosis.

34. The method of claim 30 wherein the cell proliferation disorder is polycystic
15 kidney disease.

35. The method of claim 1 wherein the composition is administered to a mammal
suffering from an infectious disease.

36. The method according to claim 33 wherein the infectious disease is selected
from the group consisting of IBD, Crohn's disease, AIDS, ARDS, and fungal diseases.

20 35. The method according to claim 1 wherein the composition is administered to a
mammal suffering from an immunological disease such as rheumatoid arthritis, autoimmune
diseases, transplant rejection, and psoriasis.

36. The therapeutically effective amount of the composition useful in the method of claim 1 wherein the composition is administered in the form of a solution.

37. The therapeutically effective amount of the composition useful in the method of claim 1 wherein the composition is administered in the form of a tablet.

5

Figure 1

Compound 187 (µg/ml)	20	10	5	0	.	.	.	0
Compound 173 (µg/ml)	.	.	.	0	20	10	5	0





Figure 2**Compound 187 (20 μ g/ml)**

Figure 3

Compound 187 (20µg/ml)
competitor oligonucleotide

-	-	+	+
-	+	-	+



INTERNATIONAL SEARCH REPORT

Intern. Application No

PCT/US 97/17013

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K38/04 A61K38/05 A61K31/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No
A	WO 92 00961 A (PFIZER) 23 January 1992 *cf. abstract, page 1, "background of the invention", claims*	1-37
A	US 5 409 944 A (BLACK W CAMERON ET AL) 25 April 1995 * cf. abstract, col. 5, lines 47-51*	1-37
A	US 5 457 237 A (DOW ROBERT L) 10 October 1995 *cf. abstract, and col. 1 bridging with col. 2, lines 1-55, col. 11, first para.*	1-37
A	WO 95 24914 A (MYOGENICS INC) 21 September 1995 *cf. abstract*	1-37

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

9 February 1998

Date of mailing of the international search report

20.02.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Stoltner, A

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 97/17013

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9200961 A	23-01-92	AT 154598 T	15-07-97
		DE 69126612 D	24-07-97
		DE 69126612 T	02-10-97
		EP 0538314 A	28-04-93
		JP 7020933 B	08-03-95
		JP 5503293 T	03-06-93
US 5409944 A	25-04-95	AU 6178894 A	26-09-94
		CA 2157107 A	14-09-94
		WO 9420480 A	15-09-94
US 5457237 A	10-10-95	EP 0586479 A	16-03-94
		JP 7119184 B	20-12-95
		JP 6502658 T	24-03-94
		WO 9221641 A	10-12-92
WO 9524914 A	21-09-95	AU 682600 B	09-10-97
		AU 2228395 A	03-10-95
		CA 2185326 A	21-09-95
		EP 0804216 A	05-11-97
		FI 963602 A	04-11-96
		JP 9511501 T	18-11-97
		NO 963814 A	14-11-96
		US 5693617 A	02-12-97

THIS PAGE BLANK (USPTO)